

Your CREXONT® (carbidopa and levodopa) extended-release capsules treatment journal



Tracking your medication and your progress

CREXONT is different from other Parkinson's medications you may have tried, and everyone's experience with CREXONT will be unique. Your dosing regimen will be individualized, and may be adjusted by your doctor based on your symptoms. This is why it's important to keep track of how you're doing on CREXONT.

How do you know if your dosing regimen needs adjusting?

You'll want to tell your healthcare provider if you're having:

- Too much "Off" time
- A delay in "On" time
- Troublesome dyskinesia

To simplify the process of tracking your response to CREXONT, we developed the CREXONT Treatment Journal. You'll find a supply of these blank journals on the following pages.

Tracking your progress with this tool is easy. Each time you take CREXONT, simply document (or have your care partner document):

- When you took it
- How much you've taken
- Your meals
- When you experience "Off" time, "On" time, dyskinesia, or other symptoms

Be sure to fill in each treatment journal page as directed above, and share them with your healthcare provider at your next check in. And remember to save this PDF to your desktop so that you can easily print additional journal pages, as needed.

Should you experience any issues, call your healthcare provider right away.

INDICATION


CREXONT (carbidopa and levodopa) extended-release capsules is a prescription medication for the treatment of Parkinson's disease, Parkinson's disease caused by infection or inflammation of the brain or Parkinson's disease-like symptoms that may result from carbon monoxide or manganese poisoning in adults.

IMPORTANT SAFETY INFORMATION

Do not take CREXONT with antidepressant medications known as nonselective monoamine oxidase (MAO) inhibitors.

Do not take CREXONT with other carbidopa-levodopa preparations without consulting your healthcare provider.

Please see additional Important Safety Information on other pages and Prescribing Information on www.CREXONT.com.



Date: _____


Complete this chart to document your daily experience with CREXONT and provide valuable insights to your healthcare provider. Multiple boxes may be checked per hour. Heres how to fill it out:

- Hourly Record: For each hour (e.g., 12:00-12:59), note details in the designated boxes.
- Dosage: Write the dosage (mg) of each CREXONT dose. If nothing was taken write N/A.
- Sleep: Mark an "X" in the sleep column for any hour you're asleep.
- On/Off Times & Dyskinesia: Place an "X" in the appropriate column if you experience "On" time, "Off" time, or Dyskinesia.
- Other Symptoms: Write down any additional symptoms as they occur.

Time (AM)	Strength of CREXONT (mg)	Meal (what you ate, if nothing write N/A)	Sleep	"Off" time	"On" time	Dyskinesia	Other symptoms
12 AM							
1 AM							
2 AM							
3 AM							
4 AM							
5 AM							
6 AM							
7 AM							
8 AM							
9 AM							
10 AM							
11 AM							

Should any questions or concerns arise as you fill out this journal, call your healthcare provider right away.

Please see additional Important Safety Information on other pages and Prescribing Information on www.CREXONT.com.



Date: _____


Complete this chart to document your daily experience with CREXONT and provide valuable insights to your healthcare provider. Multiple boxes may be checked per hour. Heres how to fill it out:

- Hourly Record: For each hour (e.g., 12:00-12:59), note details in the designated boxes.
- Dosage: Write the dosage (mg) of each CREXONT dose. If nothing was taken write N/A.
- Sleep: Mark an "X" in the sleep column for any hour you're asleep.
- On/Off Times & Dyskinesia: Place an "X" in the appropriate column if you experience "On" time, "Off" time, or Dyskinesia.
- Other Symptoms: Write down any additional symptoms as they occur.

Time (PM)	Strength of CREXONT (mg)	Meal (what you ate, if nothing write N/A)	Sleep	"Off" time	"On" time	Dyskinesia	Other symptoms
12 PM							
1 PM							
2 PM							
3 PM							
4 PM							
5 PM							
6 PM							
7 PM							
8 PM							
9 PM							
10 PM							
11 PM							

Should any questions or concerns arise as you fill out this journal, call your healthcare provider right away.

Please see additional Important Safety Information on other pages and Prescribing Information on www.CREXONT.com.



IMPORTANT SAFETY INFORMATION (continued)

CREXONT may cause falling asleep during activities of daily living, somnolence, or dizziness. Avoid activities that require alertness such as driving and operating machinery until you know how CREXONT affects you.

Your doctor should evaluate vitamin B6 levels before and during treatment with carbidopa/levodopa therapies as seizures associated with low levels of vitamin B6 have been reported. Traditional anti-seizure medications were not effective, and seizures were only resolved after vitamin B6 administration. Other symptoms of vitamin B6 deficiency include depression, confusion, cracked corners of the mouth, swollen tongue, skin inflammation, low red blood cell count, and/or numbness, tingling, or weakness in the extremities. Your doctor may advise you to take vitamin B6 as necessary.

The most common side effects that may occur with CREXONT are nausea and anxiety.

It is important to avoid sudden discontinuation or rapid dose reduction of CREXONT. If you are discontinuing CREXONT, work with your healthcare provider to taper the dose over time to reduce the risk of fever or confusion.

You may take CREXONT with or without food, but taking it with food may decrease or delay its effect. Consider taking the first dose of the day about 1 to 2 hours before eating.

Swallow CREXONT whole. Do not chew, divide, or crush the capsules.

Do not take CREXONT with alcohol.

Tell your healthcare provider if you:

- Have any heart conditions, especially if you have had a heart attack or irregular heartbeats
- Experience hallucinations or abnormal thoughts and behaviors
- Have an inability to control urges to gamble, have increased sexual urges, or experience other intense urges
- Have thoughts of suicide or have attempted suicide
- Have abnormal involuntary movements that appear or get worse during treatment
- Have ever had a peptic ulcer or glaucoma
- Become or intend to become pregnant. Based on animal data, CREXONT may cause fetal harm
- Are breastfeeding during therapy

To report SUSPECTED ADVERSE REACTIONS, contact Amneal Global Patient Safety at 1-877-835-5472, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

Please see additional Important Safety Information on other pages and Prescribing Information on www.CREXONT.com.

Amneal

CREXONT® is a registered trademark of Amneal Pharmaceuticals LLC.
© 2026 Amneal Pharmaceuticals LLC. Distributed by Amneal Specialty, a division of Amneal Pharmaceuticals LLC.
All rights reserved. PP-ADW-CREX-US-0005 05/2026

