
CREXONT®
(carbidopa and levodopa)
EXTENDED-RELEASE CAPSULES
35mg/140mg · 52.5mg/210mg
70mg/280mg · 87.5mg/350mg



STARTING YOUR TREATMENT

What to expect when starting CREXONT

A practical guide to your first weeks on CREXONT: how it works, how to take it, and how you and your doctor personalize your dose.

Please see Important Safety Information on page 11 and accompanying full Prescribing Information.

A GUIDE FOR GETTING STARTED

You've started CREXONT. Here's what comes next.

Starting a new Parkinson's medicine comes with questions. This guide walks you through what CREXONT is, how and when to take it, and what to expect as you and your doctor find the right dose for you. Remember, you are not managing this alone.

This guide will walk you through:



How CREXONT works



Finding your dose



How & when to take it

WHY THINGS CHANGE OVER TIME

Over time, your Parkinson's treatment often needs to change

Levodopa has been the standard of care treatment for Parkinson's disease. Over time, the effect of immediate-release carbidopa/levodopa (IR CD/LD) can become less predictable and symptoms can return between doses.

DID YOU KNOW?

About half of patients taking a PD medication may experience "Off" episodes within 2 to 3 years of being on treatment.

Here are some reasons your treatment could be changing:



Wearing off

Your dose may not last as long as it used to.



Ups and downs of symptom control

Through the day you may swing between stretches when your medicine is working well and your symptoms are controlled ("Good On" time), and stretches when symptoms come back — sometimes sooner than you expect ("Off" time).



Involuntary, erratic, or uncontrollable movements

Dyskinesia can appear when levodopa levels swing too high.




Delayed "On"

Constipation or slow digestion can affect how quickly your medicine starts to work.

A FAMILIAR MEDICINE, REDESIGNED TO LAST

CREXONT contains the same CD/LD you're familiar with, just redesigned to last

Discover the dual components inside CREXONT



Immediate-release granules

Dissolve quickly, so levodopa levels peak at ~1 hour


Extended-release pellets

Release levodopa slowly so levels last

Capsule and contents shown for illustrative purposes only.

The ER pellet is designed for duration*:

The ER pellets are designed to gradually release levodopa in the part of the gut where it is best absorbed*. Levodopa works best when it's absorbed in the small intestine, not the stomach.



*The exact site and duration of absorption is unknown.

Please see Important Safety Information on page 11 and accompanying full Prescribing Information.

ONE MEDICINE, FOUR STRENGTHS

CREXONT comes in 4 strengths so it can be tailored to you



35 mg / 140 mg
carbidopa / levodopa



52.5 mg / 210 mg
carbidopa / levodopa



70 mg / 280 mg
carbidopa / levodopa



87.5 mg / 350 mg
carbidopa / levodopa

Capsules shown are not actual size.

You and your doctor will work together to find which dose and frequency will help you achieve your treatment goals.

YOUR FIRST FEW WEEKS

Finding the dose that's right for you

You and your doctor will work together to personalize your dose. This takes a little time, and your feedback is the most important guide.

Day 1
Start CREXONT
2 to 4 times a day

Days 1–3
First check-in
Doctor may adjust dose and frequency

Next weeks
Fine-tuning
Some patients may need small changes

Ongoing
Your steady routine
A schedule that works for you

Safety information you should know:

What you may notice as your body adjusts

- The most common side effects are nausea and anxiety
- CREXONT may also cause sleepiness or dizziness
- Avoid driving or operating machinery until you know how it affects you


Tell your doctor right away

- Any increased "Off" time or dyskinesia
- Hallucinations or unusual thoughts or behaviors
- Strong new urges (such as gambling)
- Falling asleep during daily activities
- Thoughts of suicide

THE GOAL IS BALANCE

How your dose gets personalized

Your doctor aims for a balance: good control of your symptoms with as few side effects as possible. Small changes in timing, number of capsules, or strength can make a real difference.



What to keep track of so you can work with your doctor to get the right dose

- ☐ **How much good symptom control does your dose give you?** If your dose doesn't give you good symptom control, your doctor may need to increase your dose.
- ☐ **Are you wearing off before your next dose?** If you're wearing off between doses, your doctor may need to adjust how often you take CREXONT.
- ☐ **Are you experiencing involuntary, erratic, or uncontrollable movements?** If you're experiencing dyskinesia, your doctor may need to decrease your dose.

Be sure to talk with your doctor before changing or discontinuing any treatment. Please see Important Safety Information on page 11 and accompanying full Prescribing Information.

TAKING CREXONT EACH DAY

How and when to take CREXONT

A few simple habits help CREXONT work its best. Always follow your doctor's instructions.



3 times per day for most patients

Dosing may start at 2 to 3 times per day and may be adjusted up to 4 times per day.



Adjustments may be needed

To achieve the best balance of symptom control and tolerability, you and your doctor will find the dose and frequency that works for you.



With or without food

You can take CREXONT with or without food. A large, rich meal can make it take about 2 hours longer to work.



Swallow capsule whole

Do not chew, divide, or crush CREXONT capsules.



Sprinkle on applesauce if you have trouble swallowing

You may carefully twist apart both halves and sprinkle the entire contents on 1 to 2 tablespoons of applesauce. Consume the mixture immediately. Do not store this drug and food mixture for future use.



No alcohol

Do not take CREXONT with alcohol.

TRACK YOUR FIRST WEEKS

Your CREXONT starting log

Bring this to your appointments. Noticing patterns helps your doctor personalize your dose.

Date	Dose time	Started working	How long you had good symptom control	When symptoms returned	Notes / which symptoms

YOU'RE NOT MANAGING THIS ALONE

Questions & support to lean on

Questions to ask your doctor

- How many capsules should I take, and how many times a day?
- What should I do if my symptoms return between doses?
- Which side effects should I watch for, and which need a call?
- When should I call you versus wait until my next visit?
- What should I do if I miss a dose?

CREXONT Savings Program

Eligible commercially insured patients may qualify for a $25 co-pay on their prescriptions, applied automatically for most patients at the pharmacy.

Have Medicare Part D?

You may be able to spread your out-of-pocket costs across the year through the Medicare Prescription Payment Plan. Contact your plan to enroll in "Smoothing".

Amneal Patient Assistance Program

Offers eligible individuals the opportunity to apply to receive CREXONT free for up to one year.

My care team

Doctor / clinic _____

Phone _____

Pharmacy _____

Please see Important Safety Information on page 11 and accompanying full Prescribing Information.

Important safety information

INDICATION

CREXONT® (carbidopa and levodopa) extended-release capsules is a prescription medication for the treatment of Parkinson's disease, Parkinson's disease caused by infection or inflammation of the brain, or Parkinson's disease-like symptoms that may result from carbon monoxide or manganese poisoning in adults.

IMPORTANT SAFETY INFORMATION

- Do not take CREXONT with antidepressant medications known as nonselective monoamine oxidase (MAO) inhibitors.
- Do not take CREXONT with other carbidopa-levodopa preparations without consulting your healthcare provider.
- CREXONT may cause falling asleep during activities of daily living, somnolence, or dizziness. Avoid activities that require alertness such as driving and operating machinery until you know how CREXONT affects you.
- Your doctor should evaluate vitamin B6 levels before and during treatment with carbidopa/levodopa therapies as seizures associated with low levels of vitamin B6 have been reported. Traditional anti-seizure medications were not effective, and seizures were only resolved after vitamin B6 administration. Other symptoms of vitamin B6 deficiency include depression, confusion, cracked corners of the mouth, swollen tongue, skin inflammation, low red blood cell count, and/or numbness, tingling, or weakness in the extremities. Your doctor may advise you to take vitamin B6 as necessary.
- The most common side effects that may occur with CREXONT are nausea and anxiety.
- It is important to avoid sudden discontinuation or rapid dose reduction of CREXONT. If you are discontinuing CREXONT, work with your healthcare provider to taper the dose over time to reduce the risk of fever or confusion.
- You may take CREXONT with or without food, but taking it with food may decrease or delay its effect. Consider taking the first dose of the day about 1 to 2 hours before eating.
- Swallow CREXONT whole. Do not chew, divide, or crush CREXONT capsules. If you have difficulty swallowing the capsule, twist apart both halves and sprinkle the entire contents of both halves of the capsule on a small amount of applesauce (1 to 2 tablespoons). Swallow the mixture immediately without chewing. Do not store the drug/food mixture for future use.
- Do not take CREXONT with alcohol.

Tell your healthcare provider if you:

- Have any heart conditions, especially if you have had a heart attack or irregular heartbeats
- Experience hallucinations or abnormal thoughts and behaviors
- Have an inability to control urges to gamble, have increased sexual urges, or experience other intense urges
- Have thoughts of suicide or have attempted suicide
- Have abnormal involuntary movements that appear or get worse during treatment
- Have ever had a peptic ulcer or glaucoma
- Become or intend to become pregnant. Based on animal data, CREXONT may cause fetal harm
- Are breastfeeding during therapy

To report SUSPECTED ADVERSE REACTIONS, contact Amneal Global Patient Safety at 1-877-835-5472, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. Please read the full Prescribing Information at amneal.com/pi/crexont.pdf. For more information talk to your healthcare provider.

*"Good On" time is defined as "On" time without troublesome dyskinesia. CREXONT® is a registered trademark of Amneal Pharmaceuticals LLC. © 2026 Amneal Pharmaceuticals LLC. Distributed by Amneal Specialty, a division of Amneal Pharmaceuticals LLC. All rights reserved.

PP-PAT-CREX-US-0004 09/2026

CREXONT® (carbidopa and levodopa) EXTENDED-RELEASE CAPSULES 35mg/140mg · 52.5mg/210mg 70mg/280mg · 87.5mg/350mg